

BREYANZI is for the treatment of large B-cell lymphoma in patients when at least 2 previous treatments have not worked or have stopped working. BREYANZI is a medicine made from your own white blood cells; the cells are genetically modified to recognize and attack your lymphoma cells.



Questions to ask your doctor about Breyanzi®

About Breyanzi

What is Breyanzi?

How is Breyanzi different from treatments I've had before (such as chemotherapy, chemotherapy and immunotherapy, radiation, or stem cell transplant)?

What are the approved CAR T cell therapy options available to me?

Select Important Safety Information

What is the most important information I should know about BREYANZI?

BREYANZI may cause side effects that are life-threatening and can lead to death. Call your healthcare provider or get emergency help right away if you get any of the following:

- difficulty breathing
- fever (100.4°F/38°C or higher)
- chills/shaking chills
- confusion
- severe nausea, vomiting, diarrhea
- fast or irregular heartbeat
- dizziness/lightheadedness
- severe fatigue or weakness

It is important that you tell your healthcare providers that you have received BREYANZI and to show them your BREYANZI Patient Wallet Card. Your healthcare provider may give you other medicines to treat your side effects.

Please see Important Safety Information throughout and full Prescribing Information, including Boxed WARNINGS and Medication Guide.

Breyanzi[™]
(lisocabtagene maraleucel) SUSPENSION FOR IV INFUSION

Considering treatment with Breyanzi®

When should I start considering Breyanzi?

What factors will help you determine if Breyanzi is right for me?

Do treatments I've had affect my ability to get Breyanzi now or in the future?
What about other treatments I may be considering?

If you decide that Breyanzi is right for me, what do we need to do next?

How Breyanzi works

How does Breyanzi work with *my* immune system?

Select Important Safety Information

How will I receive BREYANZI?

- BREYANZI is made from your own white blood cells, so your blood will be collected by a process called leukapheresis.
- It takes about 3-4 weeks from the time your cells are received at the manufacturing site and are available to be shipped back to your healthcare provider, but the time may vary.
- Before you get BREYANZI, you will get 3 days of chemotherapy to prepare your body.

Please see Important Safety Information throughout and full Prescribing Information, including Boxed WARNINGS and Medication Guide.

Breyanzi[™]
(lisocabtagene maraleucl) SUSPENSION FOR IV INFUSION

Receiving Breyanzi®

What's involved in the treatment process?

How long is the entire process?

Where would I receive Breyanzi?

Will I need to stay in the hospital after receiving Breyanzi?

How long do I need to stay near the authorized treatment center after receiving Breyanzi?

Select Important Safety Information

How will I receive BREYANZI? (continued)

- When your BREYANZI is ready, your healthcare provider will give it to you through a catheter placed into your vein. BREYANZI is given as infusions of 2 different cell types.
 - You will receive infusions of one cell type, immediately followed by the other cell type.
 - The time for infusion will vary, but will usually be less than 15 minutes for each of the 2 cell types.
- During the first week, you will be monitored daily by the facility where you received your treatment.
- You should plan to stay close to this location for at least 4 weeks after getting BREYANZI. Your healthcare provider will check to see that your treatment is working and help you with any side effects that may occur.

Please see Important Safety Information throughout and full Prescribing Information, including Boxed WARNINGS and Medication Guide.

Breyanzi[™]
(lisocabtagene maraleucel) SUSPENSION FOR IV INFUSION

Benefits and risks of Breyanzi®

Were there people like me in the Breyanzi clinical trial? How did they respond to treatment?

What side effects should I be aware of?

How common is it for people to experience side effects after receiving Breyanzi?
How severe are they?

How are side effects managed?

Select Important Safety Information

How will I receive BREYANZI? (continued)

- You may be hospitalized for side effects and your healthcare provider will discharge you if your side effects are under control, and it is safe for you to leave the hospital.
- Your healthcare provider will want to do blood tests to follow your progress. It is important that you have your blood tested. If you miss an appointment, call your healthcare provider as soon as possible to reschedule.

Please see Important Safety Information throughout and full **Prescribing Information**, including **Boxed WARNINGS** and **Medication Guide**.

Breyanzi[™]
(lisocabtagene maraleucl) SUSPENSION FOR IV INFUSION

Support throughout treatment

Do I need to have a care partner throughout treatment? Why?

What support programs or organizations can help?

Additional questions or notes

At your next appointment, ask your doctor if Breyanzi® is right for you.
If you would like to learn more about how Breyanzi may help, visit [Breyanzi.com](https://www.breyanzi.com).

Select Important Safety Information

What should I avoid after receiving BREYANZI?

- Do not drive, operate heavy machinery, or do other activities that could be dangerous if you are not mentally alert, for at least 8 weeks after you get BREYANZI. This is because the treatment can cause temporary memory and coordination problems, including sleepiness, confusion, dizziness, and seizures.
- Do not donate blood, organs, tissues, or cells for transplantation.

Please see Important Safety Information throughout and full Prescribing Information, including Boxed WARNINGS and Medication Guide.



Important Safety Information

What is the most important information I should know about BREYANZI?

BREYANZI may cause side effects that are life-threatening and can lead to death. Call your healthcare provider or get emergency help right away if you get any of the following:

- difficulty breathing
- fever (100.4°F/38°C or higher)
- chills/shaking chills
- confusion
- severe nausea, vomiting, diarrhea
- fast or irregular heartbeat
- dizziness/lightheadedness
- severe fatigue or weakness

It is important that you tell your healthcare providers that you have received BREYANZI and to show them your BREYANZI Patient Wallet Card. Your healthcare provider may give you other medicines to treat your side effects.

How will I receive BREYANZI?

- BREYANZI is made from your own white blood cells, so your blood will be collected by a process called leukapheresis.
- It takes about 3-4 weeks from the time your cells are received at the manufacturing site and are available to be shipped back to your healthcare provider, but the time may vary.
- Before you get BREYANZI, you will get 3 days of chemotherapy to prepare your body.
- When your BREYANZI is ready, your healthcare provider will give it to you through a catheter placed into your vein. BREYANZI is given as infusions of 2 different cell types.
 - You will receive infusions of one cell type, immediately followed by the other cell type.
 - The time for infusion will vary, but will usually be less than 15 minutes for each of the 2 cell types.
- During the first week, you will be monitored daily by the facility where you received your treatment.
- You should plan to stay close to this location for at least 4 weeks after getting BREYANZI. Your healthcare provider will check to see that your treatment is working and help you with any side effects that may occur.
- You may be hospitalized for side effects and your healthcare provider will discharge you if your side effects are under control, and it is safe for you to leave the hospital.
- Your healthcare provider will want to do blood tests to follow your progress. It is important that you have your blood tested. If you miss an appointment, call your healthcare provider as soon as possible to reschedule.

What should I avoid after receiving BREYANZI?

- Do not drive, operate heavy machinery, or do other activities that could be dangerous if you are not

mentally alert, for at least 8 weeks after you get BREYANZI. This is because the treatment can cause temporary memory and coordination problems, including sleepiness, confusion, dizziness, and seizures.

- Do not donate blood, organs, tissues, or cells for transplantation.

What are the possible or reasonably likely side effects of BREYANZI?

The most common side effects of BREYANZI are:

- fatigue
- difficulty breathing
- fever (100.4°F/38°C or higher)
- chills/shaking chills
- confusion
- difficulty speaking or slurred speech
- severe nausea, vomiting, diarrhea
- headache
- dizziness/lightheadedness
- fast or irregular heartbeat
- swelling

BREYANZI can increase the risk of life-threatening infections that may lead to death. Tell your healthcare provider right away if you develop fever, chills, or any signs or symptoms of an infection.

BREYANZI can lower one or more types of your blood cells (red blood cells, white blood cells, or platelets). After treatment, your healthcare provider will test your blood to check for this. Tell your healthcare provider right away if you get a fever, are feeling tired, or have bruising or bleeding.

Having BREYANZI in your blood may cause a false-positive HIV test result by some commercial tests.

These are not all the possible side effects of BREYANZI. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Indication

BREYANZI is for the treatment of large B-cell lymphoma in patients when at least 2 previous treatments have not worked or have stopped working. BREYANZI is a medicine made from your own white blood cells; the cells are genetically modified to recognize and attack your lymphoma cells.

Please see full Prescribing Information, including Boxed WARNINGS and Medication Guide.


Breyanzi[™]
(lisocabtagene maraleucl) SUSPENSION FOR IV INFUSION